



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2013

---

## **Nutrition in ultra-endurance racing - aspects of energy balance, fluid balance and exercise-associated hyponatremia**

Knechtle, B

**Abstract:** Ultra-endurance athletes try to extend their limits in performance. In ultra-endurance races, athletes face limits in nutrition regarding both energy intake and fluid metabolism. The purpose of this review is to focus on the decrease in body mass, aspects of energy and fluid balance, and exercise-associated hyponatremia in ultra-endurance performance. An ultra-endurance performance lasting 24 hours or longer may lead to an energy deficit of approximately 7,000 kcal per day. This energy deficit may result in a decrease of body mass, including a decrease in both fat mass and skeletal muscle mass. The energy deficit cannot be completely compensated by increasing energy intake. Adequate fluid intake is required during an ultra-endurance performance to prevent dehydration. In case of fluid overload, both exercise-associated hyponatremia and swelling of limbs may occur. Limited fluid intake of approximately 300-400 ml per hour may prevent both exercise-associated hyponatremia and swelling of limbs. In summary, in ultra-endurance performances, an energy deficit seems to be unavoidable, and athletes are at risk to develop both exercise-associated hyponatremia and limb swelling in case of fluid overload.

DOI: <https://doi.org/10.5604/17342260.1081297>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-92140>

Journal Article

Published Version

Originally published at:

Knechtle, B (2013). Nutrition in ultra-endurance racing - aspects of energy balance, fluid balance and exercise-associated hyponatremia. *Medicina Sportiva*, 17(4):200-210.

DOI: <https://doi.org/10.5604/17342260.1081297>

# NUTRITION IN ULTRA-ENDURANCE RACING – ASPECTS OF ENERGY BALANCE, FLUID BALANCE AND EXERCISE-ASSOCIATED HYPONATREMIA

**Beat Knechtle**

Gesundheitszentrum St. Gallen, St. Gallen, Switzerland

Institute of General Practice and for Health Services Research, University of Zurich, Zurich, Switzerland

## Abstract

Ultra-endurance athletes try to extend their limits in performance. In ultra-endurance races, athletes face limits in nutrition regarding both energy intake and fluid metabolism. The purpose of this review is to focus on the decrease in body mass, aspects of energy and fluid balance, and exercise-associated hyponatremia in ultra-endurance performance. An ultra-endurance performance lasting 24 hours or longer may lead to an energy deficit of approximately 7,000 kcal per day. This energy deficit may result in a decrease of body mass, including a decrease in both fat mass and skeletal muscle mass. The energy deficit cannot be completely compensated by increasing energy intake. Adequate fluid intake is required during an ultra-endurance performance to prevent dehydration. In case of fluid overload, both exercise-associated hyponatremia and swelling of limbs may occur. Limited fluid intake of approximately 300-400 ml per hour may prevent both exercise-associated hyponatremia and swelling of limbs. In summary, in ultra-endurance performances, an energy deficit seems to be unavoidable, and athletes are at risk to develop both exercise-associated hyponatremia and limb swelling in case of fluid overload.

**Keywords:** *energy deficit, fluid overload, exercise-associated hyponatremia, limb swelling*

## Ultra-endurance performances

An ultra-endurance performance is defined as an endurance performance lasting for six hours or more [1]. The most popular ultra-endurance disciplines are swimming, cycling, running and triathlon as the combination of these three disciplines. Table 1 summarizes well-known ultra-endurance races. When ultra-endurance athletes compete for hours, days or even weeks, they face different problems regarding nutrition such as dehydration, glycogen depletion, gastrointestinal discomfort and fluid overload, which may occur as a single problem or a combination of problems. The continuous physical stress leads to an energy deficit. Furthermore, the ultra-endurance performance may lead to dehydration due to sweating.

Literature regarding nutrition in endurance and ultra-endurance has already been reviewed for specific topics [1-8]. In these reviews, recommendations were made for nutritional practices in order to recover and prepare for daily training [1,6,7], protein intake during training [6,8], intake of fat prior to an endurance performance [3], intake of carbohydrates before performance [2,4,6,8-10], intake of carbohydrates during exercise [3,4,6,10,11], intake of carbohydrates after performance [4,6,10-12], protein intake for recovery [10], and fluid intake during performance [4-8,10,11,13,14].

However, the problems of energy deficit and fluid metabolism in ultra-endurance athletes have not been

addressed. The main problems regarding energy and fluid metabolism of an ultra-endurance performance can be separated in two categories. The first aspect concerns the energy deficit with the corresponding loss in solid body masses such as fat mass and skeletal muscle mass. The second aspect concerns the deregulation of fluid metabolism with dehydration or fluid overload with an increased risk of exercise-associated hyponatremia (EAH). The aim of this review is to focus on the decrease in body mass, and the problems in energy and fluid balance ultra-endurance performance.

## Nutritional problems associated with ultra-endurance performance

### *Energy turnover and energy deficit in ultra-endurance*

Most research conducted on extreme endurance (>3 hours) is based on case studies and studies involving a small number of individuals [5]. Conducting research in these events is challenging and the number of studies is very limited. Due to the lack of research and the complexities in conducting research in ultra-endurance races, we are dependent on scientific findings from other similar endurance sports. Thus findings for nutrition in endurance performance shorter than six hours in duration are examined first. For endurance exercise lasting 30 min or more, the most likely contributors to fatigue are dehydration and carbohydrate depletion [7], whereas

gastrointestinal problems, hyperthermia, and EAH can reduce endurance exercise performance and are potentially health threatening, especially in events of four hours or longer in duration [7,15]. Gastrointestinal problems occur frequently, especially in long-distance triathlons [15]. Problems seem to be related to the intake of highly concentrated carbohydrate solutions, or hyperosmotic drinks, and the intake of fibers, fat and protein [15].

Meeting macronutrient and fluid intake demands is of highest priority for ultra-endurance athletes [4]. An ultra-endurance athlete competing for hours or days with or without breaks expends considerable amounts of energy [16-37]. Meeting the energy demands of ultra-endurance athletes during racing requires careful planning and monitoring of food and fluid intake [25,38]. Adequate carbohydrate and fluid intake during an endurance performance may help to reduce fatigue and enhance performance [10]. Current evidence continues to support mandatory high carbohydrate intakes before an event to maximize muscle glycogen stores and during an event to prevent hypoglycemia [4]. Numerous case reports [17,25,28-34,39] and field studies [19,24,27,40-44] showed, however, that ultra-endurance athletes were unable to self-regulate diet or exercise intensity to prevent a negative energy balance. Estimation of energy expenditure prior to the beginning of an ultra-endurance event would allow athletes to plan the diet energy intake better [44]. Furthermore, the

insufficient energy intake is associated with malnutrition such as a low antioxidant vitamin intake [45].

Adequate food and fluid intake is related to a successful finish of an ultra-endurance race [24,46,47]. An important key to a successful finish in an ultra-endurance race seems to be an appropriate nutrition strategy during the race [47]. Additionally, an energy deficit inhibits ultra-endurance performance. A significant negative relationship between energy intake and time taken to complete a 384-km cycle race is documented in ultra-cyclists [43]. An ultra-endurance performance results in an energy deficit [17,19-31,34,36-39,48-55]. There are reports of energy expenditures of ~365-750 kcal/hour with total energy expenditures of ~18 000-80 000 kcal, which were required to complete adventure races. These energy expenditures were accompanied by significant negative energy balances during competitions [8].

In Table 2, results from case and field studies are summarized and separated by discipline (*i.e.* swimming, cycling, running and the combination as triathlon). Regarding the single disciplines, the energy deficit seems higher in swimming compared to cycling and running. This might be explained by the different environment (*i.e.* water) as compared to cycling and running. Alternatively, swimming may involve more skeletal muscle mass than cycling or running. Or it could be that swimmers ingest fewer calories during performance than runners or cyclists do. For events lasting 24 hours or longer, the energy deficit is highest

Table 1. Well-known ultra-endurance races in swimming, cycling, running and triathlon

| Discipline                    | Distance                                | Website  |
|-------------------------------|---|--|
| Swimming                      |   |  |
| Channel Swimming              | ~34 km, sea water                       | <a href="http://www.dover.uk.com/channelswimming">www.dover.uk.com/channelswimming</a> |
| Cycling                       |   |  |
| Furnace Creek 508             | ~820 km in California, USA              | <a href="http://www.adventurecorps.com">www.adventurecorps.com</a>                     |
| Tortour                       | ~1,000 km around Switzerland            | <a href="http://www.tortour.ch">www.tortour.ch</a>                                     |
| Race around Ireland           | ~2,200 km around Ireland                | <a href="http://www.racearoundireland.com">www.racearoundireland.com</a>               |
| Race around Austria           | ~2,200 km around Austria                | <a href="http://www.racearoundaustria.at">www.racearoundaustria.at</a>                 |
| Race across America           | ~4,800 km across America                | <a href="http://www.raceacrossamerica.org">www.raceacrossamerica.org</a>               |
| Running                       |   |  |
| Ultra Trail du Mont Blanc     | 166 km                                  | <a href="http://www.ultratrailmb.com">www.ultratrailmb.com</a>                         |
| Badwater                      | 217 km                                  | <a href="http://www.badwater.com">www.badwater.com</a>                                 |
| Spartathlon                   | 246 km                                  | <a href="http://www.spartathlon.gr">www.spartathlon.gr</a>                             |
| Marathon des Sables           | ~240 km                                 | <a href="http://www.darbaroud.com">www.darbaroud.com</a>                               |
| Western States Endurance Run  | 161 km                                  | <a href="http://www.ws100.com">www.ws100.com</a>                                       |
| Trans Europe Foot Race        | ~5,100 km                               | <a href="http://www.transeurope-footrace.org">www.transeurope-footrace.org</a>         |
| Triathlon                     |   |  |
| Virgina Double Iron Triathlon | 7.6 km swim, 360 km bike, 84.4 km run   | <a href="http://www.usaultratri.com">www.usaultratri.com</a>                           |
| Virgina Triple Iron Triathlon | 11.4 km swim, 540 km bike, 126.6 km run | <a href="http://www.usaultratri.com">www.usaultratri.com</a>                           |
| Deca Iron Triathlon Mexico    | 36 km swim, 1,800 km bike, 420 km run   | <a href="http://www.multisport.com.mx">www.multisport.com.mx</a>                       |

Table 2. *Energy balance in ultra-endurance athletes in different disciplines*

| Distance and/or time            | Subjects  | Total energy intake (kcal) | Total energy expenditure (kcal) | Total energy deficit (kcal) | Energy deficit in 24 hours (kcal) | Energy deficit per hour (kcal) | Reference |
|---------------------------------|-----------|----------------------------|---------------------------------|-----------------------------|-----------------------------------|--------------------------------|-----------|
| Swimming                        |           |                            |                                 |                             |                                   |                                |           |
| 26.6 km                         | 1 male    | 2,105                      | 5,540                           | - 3,435                     |                                   | - 429                          | 30        |
| 26.6 km                         | 1 male    |                            |                                 |                             |                                   | - 500                          | 48        |
| 24-h swim                       | 1 male    | 3,900                      | 11,460                          | - 7,480                     | - 7,480                           | - 311                          | 49        |
| Mean (SD)                       |           |                            |                                 |                             |                                   | - 413±95                       |           |
| Cycling                         |           |                            |                                 |                             |                                   |                                |           |
| 12 hours indoor-cycling         | 1 male    | 2,750                      | 5,400                           | - 2,647                     |                                   | - 220                          | 34        |
| 557 km in 24 hours              | 1 male    | 5,571                      | 15,533                          | - 9,915                     | - 9,915                           | - 413                          | 39        |
| 617 km in 24 hours              | 1 male    | 10,000                     | 13,800                          | - 3,800                     | - 3,800                           | - 158                          | 28        |
| 694 km in 24 hours              | 1 male    | 10,576                     | 19,748                          | - 9,172                     | - 9,172                           | - 382                          | 25        |
| 24 hours cycling                | 6 males   | 8,450                      | 18,000                          | - 9,590                     | - 9,590                           | - 399                          | 26        |
| 1,000 km in 48 hours            | 1 male    | 12,120                     | 16,772                          | - 4,650                     | - 2,325                           | - 96                           | 36        |
| 1,126 km in 48 hours            | 1 male    | 11,098                     | 14,486                          | - 3,290                     | - 1,645                           | - 65                           | 50        |
| 2,272 km in 5 d 7 h             | 1 male    | 51,246                     | 80,800                          | - 29,554                    | - 5,585                           | - 232                          | 17        |
| 4,701 km in 9 d 16 h            | 1 male    | 96,124                     | 179,650                         | - 83,526                    | - 8,352                           | - 360                          | 22        |
| Mean (SD)                       |           |                            |                                 |                             | - 6,298<br>± 3,392                | - 258<br>± 134                 |           |
| Running                         |           |                            |                                 |                             |                                   |                                |           |
| 160 km in 20 h                  | 1 male    | 9,600                      | 8,480                           | - 1,120                     |                                   | - 56                           | 51        |
| 320 km in 54 h                  | 1 male    | 14,760                     | 18,120                          | - 3,360                     | - 1,493                           | - 62                           | 23        |
| 501 km in 6 days                | 1 male    | 39,666                     | 54,078                          | - 14,412                    | - 2,402                           | - 100                          | 21        |
| Atacama crossing                | 1 male    | 37,191                     | 101,157                         | - 63,966                    | - 3,046                           | - 127                          | 37        |
| 100 km                          | 11 female | 570                        | 6,310                           | - 5,750                     |                                   | - 452                          | 52        |
| 100 km                          | 27 male   | 760                        | 7,420                           | - 6,660                     |                                   | - 580                          | 53        |
| Mean (SD)                       |           |                            |                                 |                             | - 2,313<br>± 780                  | - 229<br>± 227                 |           |
| Triathlon                       |           |                            |                                 |                             |                                   |                                |           |
| Triple Iron ultra-triathlon     | 1 male    | 15,750                     | 27,485                          | - 11,735                    | - 6,869                           | - 286                          | 54        |
| Triple Iron ultra-triathlon     | 1 male    | 22,500                     | 28,600                          | - 6,100                     | - 3,404                           | - 141                          | 31        |
| Gigathlon multi-stage triathlon | 1 male    | 38,676                     | 59,622                          | 20,646                      | - 9,937                           | - 414                          | 29        |
| 10 x Ironman triathlon          | 1 male    | 77,640                     | 89,112                          | - 11,480                    | - 7,544                           | - 314                          | 55        |
| Mean±SD                         |           |                            |                                 |                             | - 6,938<br>± 2,699                | - 288<br>± 112                 |           |

in multi-sports disciplines and cycling. In running, the energy deficit is approximately three times lower than it is in both triathlon and cycling.

### ***Change in body mass during an ultra-endurance performance***

A further important fact in ultra-endurance racing is the finding that an ultra-endurance performance generally results in a loss in body mass (Table 3) [17,21-23,25,27,28,31,35-37,48,49,52,54,55-68]. The loss in body mass occurs most often in the lower trunk [21,37,60]. Depending on the length of the en-

durance performance and the discipline, the decrease in body mass corresponds to a decrease in fat mass [17,23,26,33,34,55,56,62-67] and/or skeletal muscle mass [17,24,32,55,56,58,59,61,62,66].

The total amount of amino acid oxidation during endurance exercise is only 1-6% of the total energy cost of exercise [69]. The decrease in skeletal muscle mass due to protein degradation is most probably very low. Concentric endurance performance, such as cycling, results in a decrease in fat mass [34,65], whereas an eccentric endurance performance, such as running, results in a decrease in muscle mass [59].

In runners, a decrease in both fat mass and skeletal muscle mass has been reported [58,59]. For swimmers, no change in body mass, fat mass or skeletal muscle mass has been reported for 12-hour indoor pool swimmers [70]. In male open-water ultra-swimmers, however, a decrease in skeletal muscle mass was observed [71].

However, also an increase in body mass can occur during an ultra-endurance performance (Table 3) [28,31,36,57,60]. Additionally, an increase in skeletal muscle mass has also been reported [28,31,34,36,54,57,60]. The increase in body mass was most probably due to fluid overload. An increase in estimated skeletal muscle mass might occur in cases where anthropometric methods were used and an increase in skin-fold thicknesses and limb circumferences was measured. Overall, ultra-endurance athletes seemed to lose approximately 0.5 kg in body mass and approximately 1.4 kg in fat mass where skeletal

muscle mass seemed to remain largely unchanged. Additionally, total body water seemed to increase by approximately 1.5l [35,36,52,54-58].

### ***Dehydration, fluid intake and fluid overload***

Dehydration refers both to hypohydration (*i.e.* dehydration induced prior to exercise) and to exercise-induced dehydration (*i.e.* dehydration that develops during exercise). The latter reduces aerobic endurance performance and results in increased body temperature, heart rate, perceived exertion, and possibly increased reliance on carbohydrates as a fuel source [72]. Most endurance athletes are concerned with dehydration during an ultra-endurance performance. Generally, ultra-endurance athletes do not meet their fluid demand during exercise [5]. It has been shown that body mass was lost during a 24-hour ultra-marathon [73]. However, body mass loss in ultra-endurance athletes seems to be

Table 3. *Change in body composition in ultra-endurance athletes in different disciplines*

| Distance and/or time                | Subjects   | Change in<br>body mass<br>(kg) | Change<br>in fat<br>mass (kg) | Change in<br>muscle mass<br>(kg) | Change in<br>body water<br>(l) | Reference |
|-------------------------------------|------------|--------------------------------|-------------------------------|----------------------------------|--------------------------------|-----------|
| Swimming                            |            |                                |                               |                                  |                                |           |
| 24-h swim                           | 1 male     | - 1.6                          | - 2.4                         | - 1.5                            | - 3.9                          | 49        |
| 12-h swim                           | 1 male     | - 1.1                          |                               | - 1.1                            |                                | 48        |
| Cycling                             |            |                                |                               |                                  |                                |           |
| 12-h indoor cycling                 | 1 male     | - 0.4                          | - 0.9                         | + 0.2                            |                                | 34        |
| 617 km in 24 hours                  | 1 male     | + 4.0                          | + 0.9                         | + 2.9                            |                                | 28        |
| 1,000 km within 48 hours            | 1 male     | + 2.5                          | - 1                           | + 0.4                            | + 1.8                          | 36        |
| 2,272 km in 5 d 7 h                 | 1 male     | - 2.0                          | - 0.79                        | - 1.21                           |                                | 17        |
| 4,701 km in 9 d 16 h                | 1 male     | - 5                            | -                             | -                                |                                | 22        |
| Running                             |            |                                |                               |                                  |                                |           |
| 12-h run                            | 1 male     | + 1.5                          | - 4.4                         | + 1.0                            | + 4.9                          | 57        |
| 320 km in 54 h                      | 1 male     | - 0.4                          | - 0.3                         | - 1.0                            |                                | 23        |
| 501 km in 6 days                    | 1 male     | - 3.0                          | - 6.8                         |                                  |                                | 21        |
| 100 km in 762 min                   | 11 females | - 1.5                          |                               |                                  | + 2.2                          | 52        |
| 100 km in 11:49 h:min               | 39 males   | -1.6                           | - 0.4                         | - 0.7                            | + 0.8                          | 56        |
| 338 km in 5 days                    | 21 males   |                                |                               | - 0.6                            |                                | 59        |
| 1,200 km in 17 days                 | 10 males   |                                | -3.9                          | - 2.0                            | + 2.3 l                        | 58        |
| Triathlon                           |            |                                |                               |                                  |                                |           |
| Triple Iron ultra-triathlon in 41 h | 1 male     | - 1.1                          | - 0.4                         | + 1.4                            | + 2.0                          | 54        |
| Triple Iron ultra-triathlon in 43 h | 1 male     | + 2.1                          | + 0.4                         | + 4.4                            |                                | 31        |
| Deca Iron ultra-triathlon           | 1 male     | + 3.2                          | + 2.4                         | + 2.4                            |                                | 60        |
| Quintuple Iron ultra-triathlon      | 1 male     | - 0.3                          | - 1.9                         |                                  | + 1.5                          | 35        |
| 10 x Ironman triathlon in 128 h     | 1 male     | - 1.0                          | - 0.8                         | - 0.9                            | + 2.8                          | 55        |
| Ironman in 11 h 36 min              | 27 males   | - 1.8                          |                               | - 1.0                            |                                | 61        |
| Triple Iron ultra-triathlon         | 31 males   | - 1.7                          | - 0.6                         | - 1.0                            |                                | 62        |
| 10 x Ironman triathlon in 128 h     | 8 males    |                                | - 3                           |                                  |                                | 63        |
| Mean±SD                             |            | - 0.4 ± 2.5                    | - 1.4 ± 2.3                   | + 0.1 ± 1.9                      | + 1.5 ± 1.30                   |           |



due to a decrease in solid mass and not dehydration [62,64,74]. Endurance athletes should attempt to minimize dehydration and limit body mass losses due to sweating to 2-3% of body mass [7]. Adequate fluid intake prevents loss in body mass [41]. However, fluid overload may lead to an increase in body mass [75] and a decrease in plasma sodium [75] with the risk of EAH [75-77].

Recent trends towards excessive fluid intake have resulted in frequent reports of hyponatremic hyperhydration in ultra-distance athletes, with a greater incidence in women than in men [4]. Fluid overload may lead to a considerable increase in body mass [75]. For example, one athlete competing in a Deca Iron ultra-triathlon with 38km of swimming, 1,800km of cycling and 422 km of running in 12 d 20 h showed an increase in body mass of 8kg within the first three days [60]. Unfortunately, energy intake, energy expenditure and fluid intake were not recorded, but the changes in skinfold thickness showed that edemas during the race had occurred. In athletes with post-race increases in body mass, increased skin-fold thicknesses and limb circumferences of the lower limb were also recorded [36,60]. In an athlete with an increase in body mass, an increase in skin-fold thicknesses at four skin-fold sites was also reported [28]. Both of these races were held in rather hot environments where most probably fluid intake was rather high.

However, in athletes with decreases in body mass, an increase in lower limb skin-fold thicknesses has also been reported [17,55,67]. In one athlete with a decrease in body mass after a Triple Iron ultra-triathlon, a considerable swelling of the feet was reported [54]. The timing in measuring skin-fold thicknesses and limb circumferences post-race is important because body mass and total body water may increase days after the race is finished [54].

The increase in body mass, skin-fold thicknesses and limb circumferences was most probably due to an increase in total body water (Table 3) [36,55,78]. An increase in total body water has been reported in ultra-endurance athletes [35,36,52,54-58,63,79,80]. The question is why both the skin-fold thicknesses and total body water increased. The increase in total body water might be due to an increase in plasma volume [35,79-82], which in turn might be due to sodium retention [79,81] due to increased activity of aldosterone [35,83]. An association between an increase in plasma volume and an increase in the potassium-to-sodium ratio in urine might suggest that the increased activity of aldosterone [84] may result in retention in both sodium and fluid during an ultra-endurance performance [53]. In a multi-stage race over seven days, total mean plasma sodium content increased and was a major factor in the increase in plasma volume [79].

Apart from these pathophysiological aspects, fluid overload might also result in an increase in limb volume. A recent study showed a relationship between changes in limb volumes and fluid intake [85]. Since neither renal function nor fluid regulating hormones were associated with changes in limb volumes, fluid overload is the most likely reason for an increase in both body mass and limb volumes. An actual study showed an association between an increased fluid intake and swelling of the feet in ultra-marathoners [86].

### ***Fluid intake and exercise-associated hyponatremia (EAH)***

Fluid overload during an endurance performance might lead to exercise-associated hyponatremia (EAH), defined as a serum sodium concentration ( $[Na^+]$ )  $<135$  mmol/l during or within 24 hours of exercise [87]. EAH was first described in the literature in 1985 by Noakes *et al.* where ultra-marathoners in South Africa with hyponatremia were thought have had 'water intoxication' [88]. Three main factors are responsible for the occurrence of EAH in endurance athletes: (i) overdrinking due to biological or psychological factors; (ii) inappropriate secretion of the antidiuretic hormone (ADH), in particular, the failure to suppress ADH-secretion in the face of an increase in total body water (TBW); and (iii) a failure to mobilize  $Na^+$  from the osmotically inactive sodium stores or alternatively inappropriate osmotic inactivation of circulating  $Na^+$  [87]. In that the mechanisms causing factors (i) and (iii) are unknown, it follows that the prevention of EAH requires that athletes be encouraged to avoid overdrinking while exercising.

EAH is the most common medical complication of ultra-distance exercise and is usually caused by excessive hypotonic fluid intake [89,90]. The main reason for developing EAH is the tendency of overdrinking during an endurance performance, by either an excessive fluid consumption [76] and/or inadequate sodium intake [91]. Subjects developing EAH during an ultra-endurance performance consumed twice as much fluid as those subjects without EAH [76]. Generally, fluid overload is reported for slower athletes [92]. However, in ultra-endurance athletes, faster athletes drink more than slower athletes but did not develop EAH [93,94].

The environmental conditions seemed to influence the prevalence of EAH. EAH was a common finding in ultra-endurance races held in extreme cold [91,95] or extreme heat [75,96]. In temperate climates, EAH was, however, relatively uncommon [83,97-110]. There seemed to be a gender difference where females seemed to be at higher risk for EAH [95]. The prevalence of EAH in ultra-marathoners [101,110,114] was, however, not higher compared to marathoners [92,111-113].

The prevalence of EAH seemed also to be dependent on the discipline (Table 4). While EAH was highly prevalent in ultra-swimming [95] and ultra-running [96], the prevalence of EAH was low [99,115] or even absent [98,100] in ultra-cyclists. In addition, the length of an ultra-endurance race seemed to increase the risk for EAH. The highest prevalence of EAH has been found in Ironman triathlons [106,108], Triple Iron ultra-triathlons [109] and ultra-marathons over 161 km [75,96].

### Nutritional aspects during ultra-endurance racing

Guidelines for nutrition strategies for ultra-endurance races are absent since there has been very little research into optimal nutritional practices in extreme sporting events. The few investigations that have indeed been conducted have mainly been case studies. No controlled studies linking specific nutritional strategies with higher performances do exist. The comprehensive nutritional analyses of the intake patterns of ultra-endurance athletes who manage to avoid health difficulties, such as gastrointestinal dis-

tress and mental disorientation commonly associated with ultra-distance efforts have yet to be compared to the nutrient and fluid intakes of ultra-athletes who fall prey to these problems. Adequate energy and fluid intake is needed to successfully compete in an ultra-endurance race [116-124]. Most studies are descriptive in nature and reporting the distribution of carbohydrates, fat and protein the athletes ingested (Table 5) [17,21,22,28,29,31,116,119,120]. Some studies report on the kind of food consumed [121-123] and some studies investigated the aspect of supplements [125-128].

### Intake of carbohydrates

Carbohydrates are the main source of energy intake for ultra-endurance athletes [17,38,55,118]. When the intake of carbohydrates, fat and protein during ultra-endurance racing was analyzed for ultra-endurance athletes, the highest percentage was found in carbohydrates. Ultra-endurance athletes consume approximately 68% of ingested energy as carbohydrates (Table 5).

Table 4. Prevalence of exercise-associated hyponatremia in ultra-endurance athletes in different disciplines

| Distance and/or time               | Conditions | Subjects                 | Prevalence of exercise-associated hyponatremia | Reference |
|------------------------------------|------------|--------------------------|--|-----------|
| Swimming                           |            |                          |  |           |
| 26-km open-water ultra-swim        | Moderate   | 25 males and 11 females  | 8 % in males and 36% in females                | 95        |
| Cycling                            |            |                          |  |           |
| 665-km mountain bike race          | Moderate   | 25 cyclists              | 0 %  | 98        |
| 109 km cycle race                  | Moderate   | 196 cyclists             | 0.5 %  | 99        |
| 720-km ultra-cycling race          | Moderate   | 65 males                 | 0 %  | 100       |
| Running                            |            |                          |  |           |
| 161-km mountain trail run          | Hot        | 45 runners               | 51 %   | 75        |
| 161-km mountain trail run          | Hot        | 47 runners               | 30 %   | 96        |
| 60-km mountain run                 | Moderate   | 123 runners              | 4 %  | 101       |
| 100-km ultra-marathon              | Moderate   | 50 male runners          | 0 %  | 83        |
| 100-km ultra-marathon              | Moderate   | 145 male runners         | 4.8 %  | 94        |
| 24-hour ultra-run                  | Moderate   | 15 males                 | 0 %  | 102       |
| 90-km ultra-marathon               | Moderate   | 626 runners              | 0.3 %  | 88        |
| 160-km trail race                  | Hot        | 13 runners               | 0 %  | 41        |
| Multi-disciplines                  |            |                          |  |           |
| 100-mile Iditasport ultra-marathon | Cold       | 8 cyclists and 8 runners | 44 %   | 91        |
| 161-km race                        | Cold       | 20 athletes              | 0 %  | 103       |
| Kayak, cycling and running         | Moderate   | 48 triathletes           | 2 %  | 104       |
| Ironman triathlon                  | Moderate   | 330 triathletes          | 1.8 %  | 105       |
| Ironman triathlon                  | Moderate   | 330 triathletes          | 18 %   | 106       |
| Ironman triathlon                  | Moderate   | 95 triathletes           | 9 %  | 107       |
| Ironman triathlon                  | Moderate   | 18 triathletes           | 28 %   | 108       |
| Triple Iron ultra-triathlon        | Moderate   | 31 triathletes           | 26 %   | 109       |

Table 5. *Intake of energy in ultra-endurance athletes in different disciplines*

| Distance and/or time        | Subjects | Intake of carbohydrates (%) | Intake of fat (%) | Intake of protein (%) | Reference |
|-----------------------------|----------|-----------------------------|-------------------|-----------------------|-----------|
| Cycling                     |          |                             |                   |                       |           |
| 617 km in 24 hours          | 1 male   | 64.2                        | 27                | 8.8                   | 28        |
| 2,272 km in 5 d 7 h         | 1 male   | 75.4                        | 14.6              | 10.0                  | 17        |
| 4,701 km in 9 d 16 h        | 1 male   | 75.2                        | 16.2              | 8.6                   | 22        |
| Running                     |          |                             |                   |                       |           |
| 100 km                      | 7 males  | 88.6                        | 6.7               | 4.7                   | 119       |
| 501 km in 6 days            | 1 male   | 40.0                        | 34.6              | 25.4                  | 21        |
| 1,005 km in 9 days          | 1 male   | 62                          | 27                | 11                    | 122       |
| Triathlon                   |          |                             |                   |                       |           |
| Deca Iron ultra-triathlon   | 1 male   | 67.4                        | 15.6              | 17.0                  | 120       |
| Gigathlon                   | 1 male   | 72                          | 14                | 13                    | 29        |
| Triple Iron ultra-triathlon | 1 male   | 72                          | 20                | 8                     | 31        |
| Mean (SD)                   |          | 68.5 ± 13.2                 | 19.5 ± 8.5        | 11.8 ± 6.1            |           |

**Intake of fat**

An increased fat intake pre-race leads to an increase in intramyocellular lipids in ultra-endurance athletes [31]. Increased intramyocellular lipids might improve ultra-endurance performance. However, field studies with controlled data do not exist with regard to whether fat loading improves ultra-endurance performance. In a case report, the ultra-endurance performance of a rower was enhanced after following a high fat diet for 14 d [129]. An increased fat intake during an ultra-endurance competition might well improve performance. However, also for this conclusion, field studies with controlled data do not exist. In a case report on an ultra-marathoner competing in a six-day ultra-marathon, the athlete consumed 34.6% fat in his daily food intake [21]. Nonetheless, body fat decreased in the first two days and was unchanged until the end of the race. In addition, performance slowed down after the first two days. Ultra-endurance athletes consume approximately 19% of energy as fat, which is higher than consumed energy in the form of protein (Table 5).

**Intake of protein**

Regarding protein intake, athletes consume approximately 12% of ingested energy as protein during racing. An observational field study at the 'Race across America' showed that ultra-endurance cyclists ingest rather large amounts of protein [121]. One might assume that athletes experienced a loss in skeletal muscle mass and tried to prevent this loss by using amino acids. A recent study tried to investigate whether an increase in amino acids during an ultra-marathon prevents skeletal muscle damage [130]. The intake of amino acids showed no effect on parameters related to skeletal muscle damage.

**Intake of ergogenic supplements, vitamins and minerals**

A dietary supplement, such as vitamins and minerals, is characterized as a product which can be used to address physiological or nutritional issues in sports. It may provide a convenient or practical means of consuming special nutrient requirements for sport activity, or it may be used to prevent/reverse nutritional deficiencies that commonly occur in athletes. The basis of the dietary supplement is an understanding of nutritional requirements and physiological effects of exercise. When the supplement is used to meet a physiological/nutritional goal triggered by sport activity, it may be proven to enhance sports performance [131]. Vitamin and mineral supplements are frequently used by competitive and recreational ultra-endurance athletes during training [122,123,126,127] and competitions [120-123].

The intake of ergogenic supplements, vitamins and minerals in ultra-endurance athletes and its effect on performance has been investigated in a number of studies [125,126,128]. In long-distance triathletes, over 60% of the athletes reported using vitamin supplements, of which vitamin C (97.5%), vitamin E (78.3%), and multivitamins (52.2%) were the most commonly used supplements during training. Almost half (47.8%) the athletes who used supplements did so to prevent or reduce cold symptoms [128]. The regular intake of vitamins and minerals seems, however, not to enhance ultra-endurance performance [125,126]. In the 'Deutschlandlauf 2006' of over 1,200 km in 17 consecutive stages, athletes with a regular intake of vitamin and mineral supplements during the four weeks before the race finished the competition no faster than athletes without vitamins



and minerals [125]. In a Triple Iron ultra-triathlon, athletes with a regular intake of vitamin and mineral supplements prior to the race were not faster [126]. Caffeine can be used as an ergogenic aid to help competitors stay awake during prolonged periods, enhance glycogen resynthesis and endurance performance [8].

### Fluid intake during endurance performance

*Ad libitum* fluid intake seemed to be the best strategy to prevent EAH and to maintain plasma sodium concentration [52,83,94,132-135]. Low fluid intake between 300 ml/h and 400 ml/h seemed to prevent EAH [52,106,132]. In a 4 h march a mean *ad libitum* fluid intake of approximately 400 ml/h maintained serum sodium concentration [132]. A fluid consumption of approximately 400 ml/h prevented from EAH in a 161-km race in the cold [103].

### Sodium supplementation during endurance performance

One might argue that a supplement of sodium during an endurance race might prevent EAH. However, two studies on Ironman triathletes showed that *ad libitum* sodium supplementation was indeed not necessary to preserve serum sodium concentrations in athletes competing for about 12 hours in an Ironman [136,137].

### Conclusions and implications for future research

Regarding these findings ultra-endurance athletes face a decrease in body mass most probably due to a decrease in both fat mass and skeletal muscle mass. During a race, athletes are not able to compensate for their energy deficit. Athletes tend to increase their fluid intake with increasing length of an ultra-endurance performance, which seems to result in both an increased risk of EAH and limb swelling. In summary, an energy deficit seems to be unavoidable in ultra-endurance performances. The best strategy to prevent both EAH and limb swelling is to minimize fluid intake to approximately 300-400 ml per hour.

### Declaration of interest

The authors report no conflicts of interest.

### References

1. Zaryski C, Smith DJ. Training principles and issues for ultra-endurance athletes. *Curr Sports Med Rep* 2005; 4: 165-70.
2. Brown RC. Nutrition for optimal performance during exercise: carbohydrate and fat. *Curr Sports Med Rep* 2002; 4: 222-9.
3. Lambert EV, Goedecke JH. The role of dietary macronutrients in optimizing endurance performance. *Curr Sports Med Rep* 2003; 4: 194-201.
4. Peters EM. Nutritional aspects in ultra-endurance exercise. *Curr Opin Clin Nutr Metab Care* 2003; 4: 427-34.
5. Rehrer NJ. Fluid and electrolyte balance in ultra-endurance sport. *Sports Med* 2001; 31: 701-15.
6. American Dietetic Association; Dietitians of Canada; American College of Sports Medicine, Rodriguez NR, Di Marco NM, Langley S. American College of Sports Medicine position stand. Nutrition and athletic performance. *Med Sci Sports Exerc* 2009; 41: 709-31.
7. Jeukendrup AE. Nutrition for endurance sports: marathon, triathlon, and road cycling. *J Sports Sci* 2011; 29: 91-9.
8. Ranchordas MK. Nutrition for adventure racing. *Sports Med* 2012; 42: 915-27.
9. Hargreaves M, Hawley JA, Jeukendrup A. Pre-exercise carbohydrate and fat ingestion: effects on metabolism and performance. *J Sports Sci* 2004; 22: 31-8.
10. Burke LM, Millet G, Tarnopolsky MA, International Association of Athletics Federations. Nutrition for distance events. *J Sports Sci* 2007; 25: 29-38.
11. Von Duvillard SP, Braun WA, Markofski M, et al. Fluids and hydration in prolonged endurance performance. *Nutrition* 2004; 7-8: 651-6.
12. Millard-Stafford M, Childers WL, Conger SA, et al. Recovery nutrition: timing and composition after endurance exercise. *Curr Sports Med Rep* 2008; 7: 193-201.
13. Maughan RJ, Shirreffs SM. Dehydration and rehydration in competitive sport. *Scand J Med Sci Sport* 2010; 3: 40-7.
14. Maughan RJ, Shirreffs SM. Nutrition for sports performance: issues and opportunities. *Proc Nutr Soc* 2012; 71: 112-9.
15. Jeukendrup AE, Jentjens RL, Moseley L. Nutritional considerations in triathlon. *Sports Med* 2005; 35: 163-81.
16. Ardigo LP, Capelli C. Energy expenditure during the LANY Footrace 2011 - a case study. *Appl Physiol Nutr Metab* 2012 Sep 12. [Epub ahead of print].
17. Bircher S, Enggist A, Jehle T, Knechtle B. Effects of an extreme endurance race on energy balance and body composition - a case report. *J Sports Sci Med* 2006; 5: 154-62.
18. Hill RJ, Davies PS. Energy expenditure during 2 wk of an ultra-endurance run around Australia. *Med Sci Sports Exerc* 2001; 33: 148-51.
19. Hulton AT, Lahart I, Williams KL, et al. Energy expenditure in the Race Across America (RAAM). *Int J Sports Med* 2010; 31: 463-7.
20. Kimber NE, Ross JJ, Mason SL, Speedy DB. Energy balance during an ironman triathlon in male and female triathletes. *Int J Sport Nutr Exerc Metab* 2002; 12: 47-62.
21. Knechtle B, Bircher S. Changes in body composition during an extreme endurance run. *Praxis (Bern 1994)* 2005; 94: 371-7.
22. Knechtle B, Enggist A, Jehle T. Energy turnover at the Race Across America (RAAM) - a case report. *Int J Sports Med* 2005; 26: 499-503.
23. Knechtle B, Knechtle P. Run across Switzerland-effect on body fat and muscle mass. *Praxis (Bern 1994)* 2007; 96: 281-6.
24. Kruseman M, Bucher S, Bovard M, et al. Nutrient intake and performance during a mountain marathon: an observational study. *Eur J Appl Physiol* 2005; 94: 151-7.
25. White JA, Ward C, Nelson H. Ergogenic demands of a 24 hour cycling event. *Br J Sports Med* 1984; 18: 165-71.
26. Peters EM. Nutritional aspects in ultra-endurance exercise. *Curr Opin Clin Nutr Metab Care* 2003; 6: 427-34.
27. Bourrilhon C, Philippe M, Chennaoui M, et al. Energy expenditure during an ultraendurance alpine climbing race. *Wilderness Environ Med* 2009; 20: 225-33.
28. Knechtle B, Knechtle P, Müller G, Zwyssig D. Energieumsatz an einem 24 Stunden Radrennen: Verhalten von Körpergewicht und Subkutanfett. *Ö J Sportmed* 2003; 33: 11-18.
29. Knechtle B, Bisig A, Schläpfer F, Zwyssig D. Energy metabolism in long-term endurance sports: a case study. *Praxis (Bern 1994)* 2003; 92: 859-64.
30. Knechtle B, Knechtle P, Heusser D. Energieumsatz bei Langstreckenschwimmen - eine Fallbeschreibung. *Ö J Sportmed* 2004; 33: 18-23.
31. Knechtle B, Zapf J, Zwyssig D, et al. Energieumsatz und Muskelstruktur bei Langzeitbelastung: eine Fallstudie. *Schweiz Z Sportmed* 2003; 51: 180-7.
32. Knechtle B, Knechtle P, Kaul R, Kohler G. Swimming for 12 hours leads to no reduction of adipose subcutaneous tissue-a case study. *Praxis (Bern 1994)* 2007; 96: 1805-10.

33. Knechtle B, Zimmermann K, Wirth A, et al. 12 hours running results in a decrease of the subcutaneous adipose tissue. *Praxis (Bern 1994)* 2007; 96: 1423-9.
34. Knechtle B, Früh HR, Knechtle P, et al. A 12 hour indoor cycling marathon leads to a measurable decrease of adipose subcutaneous tissue. *Praxis (Bern 1994)* 2007; 96: 1071-7.
35. Knechtle B, Knechtle P, Andonie JL, Kohler G. Body composition, energy, and fluid turnover in a five-day multistage ultratriathlon: a case study. *Res Sports Med* 2009; 17: 104-120.
36. Knechtle B, Knechtle P, Kohler G. The effect of 1,000 km nonstop cycling on fat mass and skeletal muscle mass. *Res Sports Med* 2011; 19: 170-85.
37. Koehler K, Huelsemann F, de Marees M, et al. Case study: simulated and real-life energy expenditure during a 3-week expedition. *Int J Sport Nutr Exerc Metab* 2011; 21: 520-6.
38. Lindeman AK. Nutrient intake of an ultraendurance cyclist. *Int J Sport Nutr* 1991; 1: 79-85.
39. Bescós R, Rodríguez FA, Iglesias X, et al. High energy deficit in an ultraendurance athlete in a 24-hour ultracycling race. *Proc (Bayl Univ Med Cent)* 2012; 25: 124-8.
40. Francescato MP, Di Prampero PE. Energy expenditure during an ultra-endurance cycling race. *J Sports Med Phys Fitness* 2002; 42: 1-7.
41. Glace BW, Murphy CA, McHugh MP. Food intake and electrolyte status of ultramarathoners competing in extreme heat. *J Am Coll Nutr* 2002; 21: 553-9.
42. Armstrong LE, Casa DJ, Emmanuel H, et al. Nutritional, physiological, and perceptual responses during a summer ultraendurance cycling event. *J Strength Cond Res* 2012; 26: 307-18.
43. Black KE, Skidmore PM, Brown RC. Energy intakes of ultraendurance cyclists during competition, an observational study. *Int J Sport Nutr Exerc Metab* 2012; 22: 19-23.
44. Rejc E, Lazzer S, Antonutto G. Energy expenditure and dietary intake of athletes during an ultraendurance event developed by hiking, cycling and mountain climbing. *J Sports Med Phys Fitness* 2010; 50: 296-302.
45. Machefer G, Groussard C, Zouhal H, et al. Nutritional and plasmatic antioxidant vitamins status of ultra endurance athletes. *J Am Coll Nutr* 2007; 26: 311-6.
46. Stuempler KJ, Hoffman MD, Weschler LB, et al. Race diet of finishers and non-finishers in a 100 mile (161 km) mountain footrace. *J Am Coll Nutr* 2011; 30: 529-35.
47. Knechtle B, Knechtle P, Rüst CA, et al. Finishers and non-finishers in the 'Swiss Cycling Marathon' to qualify for the 'Race Across America'. *J Strength Cond Res* 2011; 25: 3257-63.
48. Knechtle B, Baumann B, Knechtle P. Effect of ultra-endurance swimming on body composition—marathon swim 2006 from Rapperswil to Zurich. *Praxis (Bern 1994)* 2007; 96: 585-9.
49. Knechtle B, Knechtle P, Kohler G, Rosemann T. Does a 24-hour ultra-swimming lead to dehydration? *J Hum Sports Exerc* 2011; 6: 68-79.
50. Stewart IB, Stewart KL. Energy balance during two days of continuous stationary cycling. *J Int Soc Sports Nutr* 2007; 4: 15.
51. O'Hara WJ, Allen C, Shephard RJ, Gill JW. LaTulippe—a case study of a one hundred and sixty kilometer runner. *Br J Sports Med* 1977; 11: 83-7.
52. Knechtle B, Senn O, Imoberdorf R, et al. Maintained total body water content and serum sodium concentrations despite body mass loss in female ultra-runners drinking ad libitum during a 100 km race. *Asia Pac J Clin Nutr* 2010; 19: 83-90.
53. Knechtle B, Senn O, Imoberdorf R, et al. No fluid overload in male ultra-runners during a 100 km ultra-run. *Res Sports Med* 2011; 19: 14-27.
54. Knechtle B, Vinzent T, Kirby S, et al. The recovery phase following a Triple Iron triathlon. *J Hum Kinet* 2009; 21: 65-74.
55. Knechtle B, Knechtle P, Schück R, et al. Effects of a Deca Iron Triathlon on body composition: a case study. *Int J Sports Med* 2008; 29: 343-51.
56. Knechtle B, Wirth A, Knechtle P, Rosemann T. Increase of total body water with decrease of body mass while running 100 km nonstop—formation of edema? *Res Q Exerc Sport* 2009; 80: 593-603.
57. Knechtle B, Zimmermann K, Wirth A, et al. 12 hours running results in a decrease of the subcutaneous adipose tissue. *Praxis (Bern 1994)* 2007; 96: 1423-9.
58. Knechtle B, Duff B, Schulze I, Kohler G. A multi-stage ultra-endurance run over 1,200 km leads to a continuous accumulation of total body water. *J Sports Sci Med* 2008; 7: 357-64.
59. Knechtle B, Kohler G. Running 338 km within 5 days has no effect on body mass and body fat but reduces skeletal muscle mass - the Isarrun 2006. *J Sports Sci Med* 2007; 6: 401-7.
60. Knechtle B, Marchand, Y. Schwankungen des Körpergewichts und der Hautfaltenstärke bei einem Athleten während eines Extremausdauerwettkampfes. *Schweiz Z Sportmed* 2003; 51: 174-8.
61. Knechtle B, Baumann B, Wirth A, et al. Male ironman triathletes lose skeletal muscle mass. *Asia Pac J Clin Nutr* 2010; 19: 91-7.
62. Knechtle B, Knechtle P, Rosemann T, Oliver S. A triple Iron triathlon leads to a decrease in total body mass but not to dehydration. *Res Q Exerc Sport* 2010; 81: 319-27.
63. Knechtle B, Salas Fraire O, Andonie JL, Kohler G. Effect of a multistage ultra-endurance triathlon on body composition: World Challenge Deca Iron Triathlon 2006. *Br J Sports Med* 2008; 42: 121-5.
64. Knechtle B, Wirth A, Knechtle P, et al. Do ultra-runners in a 24-h run really dehydrate? *Ir J Med Sci* 2011; 180: 129-34.
65. Knechtle B, Wirth A, Knechtle P, Rosemann T. An ultra-cycling race leads to no decrease in skeletal muscle mass. *Int J Sports Med* 2009; 30: 163-7.
66. Knechtle B, Duff B, Amtmann G, Kohler, G. An ultratriathlon leads to a decrease of body fat and skeletal muscle mass—the Triple Iron Triathlon Austria 2006. *Res Sports Med* 2008; 16: 97-110.
67. Knechtle B, Schwanke M, Knechtle P, Kohler G. Decrease in body fat during an ultra-endurance triathlon is associated with race intensity. *Br J Sports Med* 2008; 42: 609-13.
68. Rogers G, Goodman C, Rosen C. Water budget during ultra-endurance exercise. *Med Sci Sport. Exerc* 1997; 29: 1477-81.
69. Tarnopolsky M. Protein requirements for endurance athletes. *Nutrition* 2004; 7-8: 662-8.
70. Knechtle B, Knechtle P, Kaul R, Kohler G. No change of body mass, fat mass, and skeletal muscle mass in ultraendurance swimmers after 12 hours of swimming. *Res Q Exerc Sport* 2009; 80: 62-70.
71. Weikunat T, Knechtle B, Knechtle P, et al. Body composition and hydration status changes in male and female open-water swimmers during an ultra-endurance event. *J Sports Sci* 2012; 30: 1003-13.
72. Barr SI. Effects of dehydration on exercise performance. *Can J Appl Physiol* 1999; 24: 164-72.
73. Kao WF, Shyu CL, Yang XW, et al. Athletic performance and serial weight changes during 12- and 24-hour ultra-marathons. *Clin J Sport Med* 2008; 18: 155-8.
74. Knechtle B, Knechtle P, Rosemann T, Senn O. No dehydration in mountain bike ultra-marathoners. *Clin J Sport Med* 2009; 19: 415-20.
75. Lebus DK, Casazza GA, Hoffman MD, Van Loan MD. Can changes in body mass and total body water accurately predict hyponatremia after a 161-km running race? *Clin J Sport Med* 2010; 20: 193-9.
76. Rothwell SP, Rosengren DJ, Rojek AM, et al. Exercise-associated hyponatremia on the Kokoda Trail. *Emerg Med Australas* 2011; 23: 712-6.
77. Twerenbold R, Knechtle B, Kakebeeke TH, et al. Effects of different sodium concentrations in replacement fluids during prolonged exercise in women. *Br J Sports Med* 2003; 37: 300-303.
78. Knechtle B, Morales NP, González ER, et al. Effect of a multistage ultraendurance triathlon on aldosterone, vasopressin, extracellular water and urine electrolytes. *Scott Med J* 2012; 57: 26-32.
79. Fellmann N, Ritz P, Ribeyre J, et al. Intracellular hyperhydration induced by a 7-day endurance race. *Eur J Appl Physiol Occup Physiol* 1999; 80: 353-9.

80. Mischler I, Boirie Y, Gachon P, et al. Human albumin synthesis is increased by an ultra-endurance trial. *Med Sci Sports Exerc* 2003; 35: 75-81.
81. Leiper JB, McCormick K, Robertson JD, et al. Fluid homeostasis during prolonged low-intensity walking on consecutive days. *Clin Sci (Lond)* 1988; 75: 63-70.
82. Neumayr G, Pfister R, Hoertnagl H, et al. Renal function and plasma volume following ultramarathon cycling. *Int J Sports Med* 2005; 26: 2-8.
83. Bürge J, Knechtle B, Knechtle P, et al. Maintained serum sodium in male ultra-marathoners-the role of fluid intake, vasopressin, and aldosterone in fluid and electrolyte regulation. *Horm Metab Res* 2011; 43: 646-52.
84. Keul J, Kohler B, von Glutz G, et al. Biochemical changes in a 100 km run: carbohydrates, lipids, and hormones in serum. *Eur J Appl Physiol Occup Physiol* 1981; 47: 181-9.
85. Bracher A, Knechtle B, Gnädinger M, et al. Fluid intake and changes in limb volumes in male ultra-marathoners: does fluid overload lead to peripheral oedema? *Eur J Appl Physiol* 2012; 112: 991-1003.
86. Cejka C, Knechtle B, Knechtle P, et al. An increased fluid intake leads to feet swelling in 100-km ultra-marathoners - an observational field study. *J Int Soc Sports Nutr* 2012; 9: 11.
87. Noakes TD, Sharwood K, Speedy D, et al. Three independent biological mechanisms cause exercise-associated hyponatremia: evidence from 2,135 weighed competitive athletic performances. *Proc Natl Acad Sci U S A* 2005; 102: 18550-5.
88. Noakes TD, Goodwin N, Rayner BL, et al. Water intoxication: a possible complication during endurance exercise. *Med Sci Sports Exerc* 1985; 17: 370-5.
89. Rosner MH. Exercise-associated hyponatremia. *Phys Sportsmed* 2008; 36: 55-61.
90. Rothwell SP, Rosengren DJ. Severe exercise-associated hyponatremia on the Kokoda Trail, Papua New Guinea. *Wilderness Environ Med* 2008; 19: 42-4.
91. Stuempfle KJ, Lehmann DR, Case HS, et al. Hyponatremia in a cold weather ultraendurance race. *Alaska Med* 2002; 44: 51-5.
92. Chorley J, Cianca J, Divine J. Risk factors for exercise-associated hyponatremia in non-elite marathon runners. *Clin J Sport Med* 2007; 17: 471-7.
93. Knechtle B, Knechtle P, Rosemann T. No case of exercise-associated hyponatremia in male ultra-endurance mountain bikers in the 'Swiss Bike Masters'. *Chin J Physiol* 2011; 54: 379-84.
94. Knechtle B, Knechtle P, Rosemann T. Do male 100-km ultra-marathoners overdrink? *Int J Sports Physiol Perform* 2011; 6: 195-207.
95. Wagner S, Knechtle B, Knechtle P, et al. Higher prevalence of exercise-associated hyponatremia in female than in male open-water ultra-endurance swimmers: the 'Marathon-Swim' in Lake Zurich. *Eur J Appl Physiol* 2012; 112: 1095-1106.
96. Hoffman MD, Stuempfle KJ, Rogers IR, et al. Hyponatremia in the 2009 161-km Western States Endurance Run. *Int J Sports Physiol Perform* 2012; 7: 6-10.
97. Cuthill JA, Ellis C, Inglis A. Hazards of ultra-marathon running in the Scottish Highlands: exercise-associated hyponatraemia. *Emerg Med J* 2009; 26: 906-7.
98. Schenk K, Gatterer H, Ferrari M, et al. Bike Transalp 2008: liquid intake and its effect on the body's fluid homeostasis in the course of a multistage, cross-country, MTB marathon race in the central Alps. *Clin J Sport Med* 2010; 20: 47-52.
99. Dugas JP, Noakes TD. Hyponatraemic encephalopathy despite a modest rate of fluid intake during a 109 km cycle race. *Br J Sports Med* 2005; 39: 38.
100. Rüst CA, Knechtle B, Knechtle P, Rosemann T. No case of exercise-associated hyponatraemia in top male ultra-endurance cyclists: the 'Swiss Cycling Marathon'. *Eur J Appl Physiol* 2012; 112: 689-97.
101. Page AJ, Reid SA, Speedy DB, et al. Exercise-associated hyponatremia, renal function, and nonsteroidal anti-inflammatory drug use in an ultraendurance mountain run. *Clin J Sport Med* 2007; 17: 43-8.
102. Knechtle B, Knechtle P, Rosemann T. No exercise-associated hyponatremia found in an observational field study of male ultra-marathoners participating in a 24-hour ultra-run. *Phys Sportsmed* 2010; 38: 94-100.
103. Stuempfle KJ, Lehmann DR, Case HS, et al. Change in serum sodium concentration during a cold weather ultradistance race. *Clin J Sport Med* 2003; 13: 171-5.
104. Speedy DB, Campbell R, Mulligan G, et al. Weight changes and serum sodium concentrations after an ultradistance multisport triathlon. *Clin J Sport Med* 1997; 7: 100-3.
105. Wharam PC, Speedy DB, Noakes TD, et al. NSAID use increases the risk of developing hyponatremia during an Ironman triathlon. *Med Sci Sports Exerc* 2006; 38: 618-22.
106. Speedy DB, Noakes TD, Rogers IR, et al. Hyponatremia in ultradistance triathletes. *Med Sci Sports Exerc* 1999; 31: 809-15.
107. Speedy DB, Faris JG, Hamlin M, et al. Hyponatremia and weight changes in an ultradistance triathlon. *Clin J Sport Med* 1997; 7: 180-4.
108. Speedy DB, Noakes TD, Kimber NE, et al. Fluid balance during and after an ironman triathlon. *Clin J Sport Med* 2001; 11: 44-50.
109. Rüst CA, Knechtle B, Knechtle P, Rosemann T. Higher prevalence of exercise-associated hyponatremia in Triple Iron ultra-triathletes than reported for Ironman triathletes. *Chin J Physiol* 2012; 55: 147-55.
110. Knechtle B, Gnädinger M, Knechtle P, et al. Prevalence of exercise-associated hyponatremia in male ultraendurance athletes. *Clin J Sport Med* 2011; 21: 226-32.
111. Hsieh M, Roth R, Davis DL, et al. Hyponatremia in runners requiring on-site medical treatment at a single marathon. *Med Sci Sports Exerc* 2002; 34: 185-9.
112. Kipps C, Sharma S, Pedoe DT. The incidence of exercise-associated hyponatraemia in the London marathon. *Br J Sports Med* 2011; 45: 14-19.
113. Mettler S, Rusch C, Frey WO, et al. Hyponatremia among runners in the Zurich Marathon. *Clin J Sport Med* 2008; 18: 344-9.
114. Knechtle B, Knechtle P, Rosemann T. Low prevalence of exercise-associated hyponatremia in male 100 km ultra-marathon runners in Switzerland. *Eur J Appl Physiol* 2011; 111: 1007-16.
115. Hew-Butler T, Dugas JP, Noakes TD, Verbalis JG. Changes in plasma arginine vasopressin concentrations in cyclists participating in a 109-km cycle race. *Br J Sports Med* 2010; 44: 594-7.
116. Eden BD, Abernethy PJ. Nutritional intake during an ultraendurance running race. *Int J Sport Nutr* 1994; 4: 166-74.
117. Bescós R, Rodríguez FA, Iglesias X, et al. Nutritional behavior of cyclists during a 24-hour team relay race: a field study report. *J Int Soc Sports Nutr* 2012; 9: 3.
118. García-Rovés PM, Terrados N, Fernández SF, Patterson AM. Macronutrients intake of top level cyclists during continuous competition-change in the feeding pattern. *Int J Sports Med* 1998; 19: 61-7.
119. Fallon KE, Broad E, Thompson MW, Reull PA. Nutritional and fluid intake in a 100-km ultramarathon. *Int J Sport Nutr* 1998; 8: 24-35.
120. Knechtle B, Müller G. Ernährung bei einem Extremausdauerwettkampf. *Deut Z Sportmed* 2002; 53: 54-7.
121. Knechtle B, Pitre J, Chandler C. Food habits and use of supplements in extreme endurance cyclists - the Race Across America (RAAM) 2006. *Schweiz Z Sportmed* 2007; 55: 102-6.
122. Knechtle B, Knechtle P, Kaul R. Nutritional practices of extreme endurance swimmers - the Marathon swim in the lake of Zurich 2006. *Pak J Nutr* 2007; 6: 188-93.
123. Knechtle B, Schulze I. Nutritional behaviours in ultra-endurance runners - Deutschlandlauf 2006. *Praxis (Bern 1994)* 2008; 97: 243-51.
124. Enqvist JK, Mattsson CM, Johansson PH, et al. Energy turnover during 24 hours and 6 days of adventure racing. *J Sports Sci* 2010; 28: 947-55.
125. Knechtle B, Knechtle P, Schulze I, Kohler G. Vitamins, minerals and race performance in ultra-endurance runners-Deutschlandlauf 2006. *Asia Pac J Clin Nutr* 2008; 17: 194-8.
126. Frohnauer A, Schwanke M, Kulow W, Kohler G, Knechtle B, et al. Vitamins, minerals and race performance in ultra-endurance triathletes - Triple Iron Triathlon Lensahn 2006. *Pak J Nutr* 2008; 7: 283-6.
127. Haymes EM. Vitamin and mineral supplementation to athletes. *Int J Sport Nutr* 1991; 1: 146-69.



128. Knez WL, Peake JM. The prevalence of vitamin supplementation in ultraendurance triathletes. *Int J Sport Nutr Exerc Metab* 2010; 20: 507-14.
129. Robins AL, Davies DM, Jones GE. The effect of nutritional manipulation on ultra-endurance performance: a case study. *Res Sports Med* 2005; 13: 199-215.
130. Knechtle B, Knechtle P, Mrazek C, et al. No effect of short-term amino acid supplementation on variables related to skeletal muscle damage in 100 km ultra-runners – a randomized controlled trial. *J Int Soc Sports Nutr* 2011; 8: 6.
131. Burke LM, Read RS. Dietary supplements in sport. *Sports Med* 1993; 15: 43-65.
132. Nolte H, Noakes TD, Van Vuuren B. Ad libitum fluid replacement in military personnel during a 4-h route march. *Med Sci Sports Exerc* 2010; 42: 1675-80.
133. Nolte HW, Noakes TD, van Vuuren B. Protection of total body water content and absence of hyperthermia despite 2% body mass loss ('voluntary dehydration') in soldiers drinking ad libitum during prolonged exercise in cool environmental conditions. *Br J Sports Med* 2011; 45: 1106-12.
134. Nolte HW, Noakes TD, Van Vuuren B. Trained humans can exercise safely in extreme dry heat when drinking water ad libitum. *J Sports Sci* 2011; 29: 1233-41.
135. Tam N, Nolte HW, Noakes TD. Changes in total body water content during running races of 21.1 km and 56 km in athletes drinking ad libitum. *Clin J Sport Med* 2011; 21: 218-25.
136. Hew-Butler TD, Sharwood K, Collins M, et al. Sodium supplementation is not required to maintain serum sodium concentrations during an Ironman triathlon. *Br J Sports Med* 2006; 40: 255-9.
137. Speedy DB, Thompson JM, Rodgers I, et al. Oral salt supplementation during ultradistance exercise. *Clin J Sport Med* 2002; 12: 279-84.

Accepted: November 28, 2013

Published: December 20, 2013

Address for correspondence:

PD Dr. med. Beat Knechtle

Facharzt FMH für Allgemeinmedizin

Gesundheitszentrum St. Gallen

Vadianstrasse 26

9001 St. Gallen

Switzerland

Telephone: +41 (0) 71 226 82 82

Fax: +41 (0) 71 226 82 72

Email: beat.knechtle@hispeed.ch